



The importance of a
reliable dressing


 **smith&nephew**
DURA FIBER[®] Ag
Silver Gelling Fibre Dressing

Supporting healthcare professionals

DURAFIBER[◇] Ag

Strength in performance and design

DURAFIBER[◇] Ag is a highly absorbent^{1,2} silver gelling fibre dressing designed to provide clinicians with a versatile and reliable solution for managing infected chronic and acute wounds.



Indications

DURAFIBER[◇] Ag is designed to be left in place for up to 7 days^{2,4}
It can be applied to chronic and acute wounds including:

- Leg ulcers
- Pressure injuries
- Diabetes-related foot ulcers
- Surgical wounds
- Traumatic wounds
- Tunnelling wounds
- Partial thickness burns
- Wounds prone to bleeding following debridement



DURAFIBER[◇] Ag range

Some wounds require a dressing that is effective, easy to apply, and comfortable to the patient

Designed with patients in mind

Fast and highly absorptive properties¹²



Absorbs¹² and retains³ fluid even under compression



Locks exudate^{13,6} and any pathogenic bacteria⁷ into the dressing



Silver technology with antimicrobial activity^{8,10}



Minimal Shrinkage¹² to conform intimately¹¹ to the wound bed^{3,12}

Designed for optimal patient comfort³



Comfortable to the patient³



Fast gelling¹² provides a cool feeling when in contact with fluid³



Easy to apply³



Intact one piece removal thanks to its high wet tensile strength¹⁴



Did you know?

Our products can be used in conjunction with a secondary dressing such as ALLEVYN[®]^{14,15}




Fast acting

DURAFIBER[®] Ag has been demonstrated to start killing pathogens within 30 minutes³ of contact and to possess an antimicrobial activity, effective for up to 7 days.⁸

Silver gelling mode of action


When DURAFIBER[®] Ag comes into contact with exudate,^{13,6} the individual fibres hydrate and swell as they absorb wound fluid, forming a soft, cohesive gel sheet.¹² As the fibres swell they lock in the absorbed exudate⁶ and the silver-containing gelled dressing matrix provides an antimicrobial action proven (in vitro) effective against a broad spectrum of bacteria including Methicillin-resistant *Staphylococcus aureus* (MRSA) and *Pseudomonas aeruginosa*.⁸⁻¹⁰



1 Before application



2 Forming a gel upon contact with exudate³



3 Locks in exudate^{3,6} and harmful bacteria⁷ it may contain

Ordering information

Item Code	Size	Units per carton
66800578	5cm x 5cm	Box/10
66800579	10cm x 10cm	Box/10
66800580	15cm x 15cm	Box/5
66800581	20cm x 30cm	Box/5
66800582	2cm x 45cm	Box/5
66800583	4cm x 10cm	Box/5
66800584	4cm x 20cm	Box/5
66800585	4cm x 30cm	Box/5

DURAFIBER® Ag

Antimicrobial action⁸ with high absorption¹⁻² and strength¹⁻⁴ for optimal patient comfort.³



Smith & Nephew Pty Ltd
Australia
85 Waterloo Road
North Ryde NSW 2113
PO Box 393
North Ryde NSW 1670
T +61 2 9857 3999
F +61 2 9857 3900
www.smith-nephew.com/australia
Customer Service
T 13 13 60
F 1800 671 000

Smith & Nephew Ltd
New Zealand
Unit A 36 Hillside Road
Wairau Valley Auckland 0627
PO Box 316005
Wairau Valley Auckland 0760
T +64 9 820 2840
F +64 9 820 2841
www.smith-nephew.com/new-zealand
Customer Service
T 0800 657 799 (Surgical)
T 0800 807 663 (Wound)

®Trademark of Smith & Nephew
SN13943 (05/18)

Please consult product labels and package insert for indications, contraindications, hazards, warnings, cautions and instructions for use.

Supporting healthcare professionals
for over 150 years

1. Giaffar L. Analytical and physical properties of DURAFIBER Ag recipe E. Data on file; 2016: report DS/16/341/R. 2. Dowler A. DURAFIBER Ag (Recipe E) Testing. Data on file; 2015: report DS/15/269/R. 3. Forlee M, et al. A prospective, open, multicentre study to evaluate a new gelling fibre dressing containing silver in the management of venous leg ulcers. *Int. Wound J* 2014; 11:438-445. 4. Dowler A. Integrity testing of DURAFIBER Ag in horse serum and ionic solution. Data on file; 2012: report DS/12/093/DOF. 5. Giaffar L. DURAFIBER Ag recipe E absorption during compression. Data on file; 2016: report DS/16/443/R. 6. Lidster C. Sequestration testing of DURAFIBER Ag. Data on file; 2014: report DS/12/158/R2. 7. Lumb H. Visual demonstration of sequestration of *Pseudomonas aeruginosa* and *Staphylococcus aureus* in a DURAFIBER® Ag dressing using confocal microscopy. Data on file; 2014: report 1412017. 8. Lumb H. Antimicrobial activity of DURAFIBER® Ag against bacteria, yeast and fungi commonly found in wounds over a 7 day period. Data on file; 2015: report 1510009. 9. Lumb H. Antimicrobial activity of DURAFIBER® Ag against a broad spectrum of wound pathogens. Data on file; 2015: report 1510008. 10. Lumb H. Antimicrobial activity of DURAFIBER® Ag against bacteria, yeast and fungi commonly found in wounds over a 4 hour period. Data on file; 2015: report 15010010. 11. Lidster C. Intimate contact testing of DURAFIBER Ag. Data on file; 2014: report DS/12/143/R2. 12. Barrett S, et al. Case series evaluation: the use of DURAFIBER on exuding wounds. *Wounds UK* 2012; 8: 104-113. 13. Lumb H. Assessment of the contact kill activity of DURAFIBER Ag against common wound pathogens. Data on file; 2016: report 1609013. 14. Lidster C. Wound model testing of DURAFIBER Ag with various secondary dressings. Data on file; 2014: report DS/10/053/R2. 15. Blackburn I. DURAFIBER Ag can be used in conjunction with various secondary retentive dressing (ALLEVYN Life, ALLEVYN Non Adhesive, ALLEVYN Gentle Border, ALLEVYN Life and OPSITE). Data on file; 2014: Technical Justification Summary.