



Effective wound debridement,
rehydration and soothing¹

 **smith&nephew**
INTRASITE◊
Hydrogel Wound Dressing

INTRASITE[◊] Gel

Easy squeeze APPLIPAK[◊]

- Shaped to be easy to hold and depress
- Provides controlled delivery of gel for less mess and waste
- Saves waste by dispensing the full amount of gel

Effective debridement and desloughing¹

- By its gentle, yet effective, rehydrating action, INTRASITE can produce rapid debridement of necrotic tissue. It loosens and absorbs slough, as well as absorbs excess exudate.
- Cleans the wound to allow healing to progress¹

INTRASITE[◊] Conformable

Conformable hydrogel dressing²

- For packing wounds and awkward to dress areas, INTRASITE Gel is available in a pre-loaded non-woven dressing – INTRASITE Conformable.



Gentle action²

- Leaves delicate granulation tissue undisturbed

Bacteriostatic effect³

- Contains propylene glycol which has bacteriostatic properties


Moist wound healing

- Creates and maintains a moist wound healing environment by increasing moisture content at the wound interface

Clinically proven²

- INTRASITE is a clinically proven hydrogel that can be used on:
 - Pressure injuries
 - Leg ulcers
 - Diabetes-related foot ulcers
 - Amputation wounds
 - Extravasation
 - Surgical wounds
 - Fungating wounds
 - Burns

Pressure injury on newborn baby²



Before INTRASITE Conformable



After INTRASITE Conformable

Ordering information

If you need 8g gel choose:

Code	Product description	Items per unit
7308	8g Applipak	Box/10
66000324	10cm x 10cm	Box/10

*Not available in New Zealand

If you need 15g gel choose:

Code	Product description	Items per unit
36100614	15g Applipak	Box/10
36361354	10cm x 20cm	Box/10

If you need 25g gel choose:

Code	Product description	Items per unit
7313	25g Applipak	Box/10
66000326	10cm x 40cm	Box/10*

N.B. To avoid the dressing drying out, the size of INTRASITE Conformable needs to be selected based on the amount of gel required for the wound rather than the size of the dressing.

References

1. Flanagan M. The efficacy of a hydrogel in the treatment of wounds with non-viable tissue. *J Wound Care* 1995; 4(6): 264-267.
2. Vernon T. Intrasite Gel and Intrasite Conformable: the hydrogel range. *Brit J Nurs* 2000; 9: 1083-1088.
3. Harrow J. A comparison of the effects of two hydrogels upon the proliferation of three micro-organisms important in chronic wound care. Data on file; 2006: report 3967/2.

Smith & Nephew Pty Ltd
Australia
 T +61 2 9857 3999 F +61 2 9857 3900
www.smith-nephew.com/australia
Customer Service
 T 13 13 60 F 1800 671 000

Smith & Nephew Ltd
New Zealand
 T +64 9 820 2840 F +64 9 820 2841
www.smith-nephew.com/new-zealand
Customer Service
 T 0800 657 799 (Surgical) T 0800 807 663 (Wound)

Please consult product labels and inserts for any indications, contraindications, hazards, warnings, cautions and instructions for use.

[◊]Trademark of Smith & Nephew
 SNI2761 (11/16)